

3

ΝΟΕΜΒΡΙΟΥ

2018

ΑΘΗΝΑ

ΛΑΕΔ

Λέσχη Αξιωματικών
Ενόπλων Δυνάμεων

ΕΛΛΗΝΙΚΗ ΕΤΑΙΡΕΙΑ ΕΠΙΣΤΗΜΟΝΩΝ-ΕΡΓΑΖΟΜΕΝΩΝ
ΑΚΤΙΝΟΘΕΡΑΠΕΙΑΣ (Ε.Ε.Ε.Α.)



Σε συνεργασία με: Παθολογικό Ογκολογικό Τμήμα ΕΑΝΠ ΜΕΤΑΣ

Ισχύς εν τη ενώσει Συνδυασμένη οντιμετώπιση του καρκίνου



η θέση της ακτινοθεραπείας
στον καρκίνο του προστάτη

Ιωάννης Γεωργακόπουλος
Ακτινοθεραπευτής Ογκολόγος

δεδομένα

- τεκμηριωμένος ρόλος ακτινοθεραπείας
- κλιμάκωση της δόσης σχετίζεται με καλύτερο θεραπευτικό αποτέλεσμα
- υπεροχή IMRT/IGRT έναντι συμβατικής ακτινοθεραπείας
- συνεργικός ρόλος ακτινοθεραπείας με ορμονοθεραπεία

νεότερα δεδομένα

- υποκλασματοποίηση δόσης
- σωματιδιακές ακτινοβολίες, πρωτόνια
- χημειοθεραπεία
- μετεγχειρητική ακτινοθεραπεία, απάντηση στο κλασικό ερώτημα “επικουρική ή θεραπεία διάσωσης”;
- μελλοντικές προκλήσεις

δεδομένα

ρόλος ακτινοθεραπείας


10-Year Outcome

F.C. Hamdy, J.L. Donovan, J.
R.M. Martin, J. Oxley, M. Rot
R. Kockelbergh, H. Kyn

Patient-Reported Outcomes after Monitoring, Surgery, or Radiotherapy for Prostate Cancer

J.L. Donovan, F.C. Hamdy, J.A. Lane, M. Mason, C. Metcalfe, E. Walsh,
J.M. Blazeby, T.J. Peters, P. Holding, S. Bonnington, T. Lennon, L. Bradshaw,
D. Cooper, P. Herbert, J. Howson, A. Jones, N. Lyons, E. Salter, P. Thompson,
S. Tidball, J. Blaikie, C. Gray, P. Bollina, J. Catto, A. Doble, A. Doherty, D. Gillatt,
R. Kockelbergh, H. Kynaston, A. Paul, P. Powell, S. Prescott, D.J. Rosario, E. Rowe,
M. Davis, E.L. Turner, R.M. Martin, and D.E. Neal, for the ProtecT Study Group*

ProtecT



	Active monitoring (n=545)	Radiotherapy (n=545)	Radical prostatectomy (n=553)
Age at invitation (years)			
49-54	58 (11%)	62 (11%)	69 (12%)
55-59	140 (26%)	141 (26%)	137 (25%)
60-64	184 (34%)	176 (32%)	172 (31%)
65-69	163 (30%)	166 (30%)	175 (32%)
Median age (range)	62 (50-69)	62 (49-69)*	62 (50-69)
PSA (µg/L)			
3·0-5·9	373 (68%)	373 (68%)	371 (67%)
6·0-9·9	116 (21%)	121 (22%)	123 (22%)
≥10·0	56 (10%)	51 (9%)	59 (11%)
Median PSA (range; µg/L)	4·6 (3·0-20·9)†	4·6 (3·0-18·8)	4·7 (3·0-18·4)
Gleason score			
6	421 (77%)	423 (78%)	422 (76%)
7	111 (20%)	108 (20%)	120 (22%)
8-10	13 (2%)	14 (3%)	10 (2%)
Missing	0	0	1 (<1%)
Clinical stage			
T1c	410 (75%)	429 (79%)	410 (74%)
T2	135 (25%)	116 (21%)	143 (26%)

ProtecT

- 545 active monitoring
 - PSA q3 m x 1y, q6 m thereafter, rise of 50% in 12 m: consider biopsy
- 553 radical prostatectomy
- 545 radical radiotherapy
 - RT: 3DCRT 74 Gy (37 fr) + NAD (3-6 m)
- 10 y follow up
- Disease Progression, Metastasis, Cancer Specific Survival, Overall Survival
- Toxicity
- Quality of Life
 - baseline, 6 m, 12m and annually thereafter

ICIQ: International Consultation of Incontinence Questionnaire

EPIC: Expanded Prostate Cancer Index Composite

ICSmaleSF: International Continence Society male Short Form


SF12: Medical Outcomes Study 12

HADS: Hospital Anxiety and Depression Scale

EORTC- QLQ C30


ProtecT αποτελέσματα θεραπείας

- Disease Progression
 - AM 112 men; 22.9 per 1000 person-years; 95% CI, 19.0 to 27.5
 - RP 46 men; 8.9 per 1000 person-years; 95% CI, 6.7 to 11.9
 - RT 46 men; 9.0 per 1000 person-years; 95% CI, 6.7 to 12.0
- p<0.001
- Metastasis
 - AM 33 men; 6.3 per 1000 person-years; 95% CI, 4.5 to 8.8
 - RP 13 men; 2.4 per 1000 person-years; 95% CI, 1.4 to 4.2
 - RT 16 men; 3.0 per 1000 person-years; 95% CI, 1.9 to 4.9
- p=0.004
- Prostate Cancer Specific Survival - no significant difference




ProtecT τοξικότητα θεραπείας


EPIC Item: ≥ 1 Pad per Day



EPIC Problem with Erectile Dysfunction



EPIC Bowel Bother Score




- prostatectomy greatest negative effect on **urinary continence** at 6 months; the effect of urinary incontinence on QoL worse in the prostatectomy group for 2 years, but then similar to that reported in the other groups
- **nocturia** increased in all groups; the increase particularly in the radiotherapy group at 6 months
- at baseline 67% of men reported **erections** firm enough for intercourse, but by 6 months this rate fell to 52% in the active-monitoring group, to 22% in the radiotherapy group, and to 12% in the prostatectomy group
- **bowel function** worse in the radiotherapy group than in other groups. However, there was then considerable recovery in the radiotherapy group for these measures, apart from more frequent **bloody stools**

Hamdy, *Lancet* 2016

Donovan, *Lancet* 2016

ProtecT σχετική με την υγεία ποιότητα ζωής



Hamdy, *Lancet* 2016
Donovan, *Lancet* 2016